

Table 2 - continued Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																						
			44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	
Female	Control	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20		
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Corn Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	0.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	2.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		Corn Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	12.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Soiled perinaris		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87
Female	Control	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	0.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
		No abnormality	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	19	19
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	
Soiled perinearis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109
Female	Control	Number of examined	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	0.5 mg/kg	Number of examined	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	20	20	20	20	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
No abnormality		19	19	19	19	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Soiled perianis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																						
			110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10		
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Loss of hair	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Soiled perianis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Torticollis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Lateral position	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	10	10	10	10	10	10	10	10	10	9	9	9	9	9	9	9	9	10	9	9	9	9
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	0.5 mg/kg	Mass	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1
		Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
	2.5 mg/kg	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
	12.5 mg/kg	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Loss of hair	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soiled perianaris		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	0.5 mg/kg	Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
		Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Loss of hair	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Soiled perianis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Torticollis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Lateral position	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
No abnormality		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soiled perinearis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																				
			198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
No abnormality		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Soiled perianis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																				
			220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
No abnormality		10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Soiled perinearis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	8	8	8	8	8	8	8	8	8	8	8
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Mass	1	1	1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	2	2	
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soiled perinearis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	8	8	8	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	0.5 mg/kg	Mass	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
		Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
12.5 mg/kg	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Loss of hair	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Soiled perianis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Torticollis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Lateral position	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

Table 2 - continued Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Corn		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soiled perinaris		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																					
			308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	Mass	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	12.5 mg/kg	Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Loss of hair	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soiled perinaris		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days																						
			330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	7	7	7	7	7	7	7	7	10	7	7	7	10	10	10	10	10	10	10	10	10	10	
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Mass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	10	10	10	10	9	9	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	
		Mass	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
	12.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
		No abnormality	10	10	10	10	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	
		Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mass		0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Hypoactivity		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Soiled perianis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Torticollis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Bradypnea		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Lateral position		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Dyspnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			

Table 2 - continued

Incidence of clinical signs
Female

Study No. P030097

Sex	Group and dose	Clinical sign	Days												
			352	353	354	355	356	357	358	359	360	361	362	363	364
Female	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	7	7	7	7	7	7	7	7	7	7	7	7	7
		Abnormality of tooth	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	1	1	1	1	1	1	1	1	1	1	1	1	1
	Mass	2	2	2	2	2	2	2	2	2	2	2	2	2	
	0.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	9	9	9	9	9	9	9	9	9	9	9	9	10
		Corn	1	1	1	1	1	1	1	1	1	1	1	1	0
	2.5 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	8	8	8	8	8	8	8	8	8	8	8	8	7
		Corn	1	1	1	2	1	1	1	1	1	1	1	1	2
		Mass	1	1	1	1	1	1	1	1	1	1	1	1	1
	12.5 mg/kg	Number of examined	10	10	10	9	9	9	9	9	9	9	9	9	9
No abnormality		7	7	7	7	7	7	7	6	6	6	6	6	5	
Corn		1	1	1	1	1	1	1	1	1	1	1	1	2	
Mass		1	1	1	1	1	1	1	2	2	2	2	2	3	
Loss of hair		0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypoactivity		1	1	0	0	0	0	0	0	0	0	0	0	0	
Soiled perianaris		1	1	1	0	0	0	0	0	0	0	0	0	0	
Torticollis		1	1	1	0	0	0	0	0	0	0	0	0	0	
Bradypnea		1	1	0	0	0	0	0	0	0	0	0	0	0	
Lateral position		0	0	1	0	0	0	0	0	0	0	0	0	0	
Dyspnea		0	0	1	0	0	0	0	0	0	0	0	0	0	
Imminently sacrifice		0	0	1	0	0	0	0	0	0	0	0	0	0	

Table 3 Body weights
Male, Female

Study No. P030097

Sex	Group and dose		Body weight (g) on day										
			1	8	15	22	29	36	43	50	57	64	
Male	Control	N	20	20	20	20	20	20	20	20	20	20	20
		Mean	212.1	267.5	320.9	362.2	400.2	431.3	456.1	479.2	501.7	520.9	
	S.D.	±8.5	±14.3	±23.7	±30.2	±36.7	±40.0	±42.3	±44.9	±50.0	±52.9		
	0.1 mg/kg	N	20	20	20	20	20	20	20	20	20	20	20
		Mean	212.4	271.5	327.0	372.7	414.5	445.6	469.9	495.1	521.6	537.6	
	S.D.	±9.1	±13.6	±19.0	±24.8	±27.5	±29.6	±31.0	±33.4	±34.5	±37.3		
	0.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20	20
		Mean	211.5	274.5	334.9	381.2	423.2	456.6	481.6	505.5	526.6	545.0	
	S.D.	±9.5	±14.6	±20.3	±25.3	±30.6	±33.6	±36.0	±38.4	±40.5	±44.1		
	2.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20	20
		Mean	212.0	268.5	317.5	353.9	382.1	404.4*	421.1**	438.8**	454.4**	466.2**	
	S.D.	±10.1	±13.3	±18.7	±23.1	±25.9	±29.6	±32.2	±38.8	±42.3	±46.6		
Female	Control	N	20	20	20	20	20	20	20	20	20	20	
		Mean	157.7	187.3	210.8	235.1	249.1	261.3	268.6	276.1	286.1	290.9	
	S.D.	±8.6	±11.1	±12.6	±15.7	±16.5	±17.3	±18.1	±17.7	±21.4	±22.8		
	0.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20	
		Mean	159.1	187.9	212.3	237.2	250.9	261.9	270.2	281.2	289.5	294.3	
	S.D.	±8.3	±11.6	±13.4	±18.0	±19.3	±21.3	±21.6	±23.7	±25.5	±27.0		
	2.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20	
		Mean	159.0	190.0	211.9	238.4	253.4	265.7	273.3	284.9	293.5	298.6	
	S.D.	±7.9	±10.9	±13.1	±15.7	±17.4	±18.9	±19.8	±21.8	±23.6	±24.6		
	12.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20	
		Mean	157.1	189.1	213.8	238.3	252.7	265.5	272.6	281.5	290.2	295.1	
	S.D.	±8.4	±11.8	±13.4	±15.9	±18.3	±19.3	±18.3	±21.1	±22.5	±23.2		

*: P<0.05, **: P<0.01 (significantly different from control).

001
69

Table 3 - continued
 Body weights
 Male, Female

Study No. P030097

Sex	Group and dose		Body weight (g) on day									
			71	78	85	91	92	120	148	176	204	232
Male	Control	N	20	20	20	20	10	10	10	10	10	10
		Mean	531.7	546.4	558.8	568.3	583.0	625.2	657.5	683.9	713.3	735.3
		S.D.	±54.6	±55.4	±59.0	±60.7	±79.8	±88.8	±102.5	±111.5	±118.1	±113.7
	0.1 mg/kg	N	20	20	20	20	10	10	10	10	10	9
		Mean	553.0	569.9	583.3	589.4	590.4	616.0	643.8	668.9	693.1	730.0
		S.D.	±35.9	±38.5	±40.1	±40.0	±39.5	±64.6	±69.7	±74.8	±76.4	±74.4
	0.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	560.3	575.9	586.4	594.3	617.0	648.9	685.1	706.5	736.8	759.3
		S.D.	±47.1	±48.6	±49.8	±52.4	±52.7	±61.7	±71.0	±82.7	±90.7	±98.4
	2.5 mg/kg	N	19	19	19	19	10	10	10	10	10	10
		Mean	475.4**	483.8**	491.5**	493.2**	515.0*	535.9*	554.5*	565.4*	588.7*	593.7**
		S.D.	±46.4	±49.4	±50.8	±51.8	±61.0	±59.2	±65.0	±68.2	±74.8	±82.7
Female	Control	N	20	20	20	20	10	10	10	10	10	
		Mean	296.2	302.5	309.1	315.2	308.3	321.7	334.9	348.6	359.3	375.7
		S.D.	±22.5	±24.6	±25.7	±26.8	±26.9	±29.4	±35.8	±41.2	±47.1	±56.3
	0.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	299.0	307.4	312.2	314.0	309.7	325.2	341.8	354.4	365.1	380.6
		S.D.	±27.1	±28.5	±29.7	±30.6	±28.5	±36.2	±36.1	±41.0	±42.4	±46.3
	2.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	301.9	311.4	316.8	320.4	328.9	350.7	362.8	376.5	391.4	411.8
		S.D.	±26.0	±28.0	±30.3	±31.7	±39.3	±46.0	±49.2	±54.3	±60.6	±70.8
	12.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	300.3	306.0	309.1	311.9	307.5	324.0	337.7	348.3	361.7	373.7
		S.D.	±23.6	±24.8	±27.4	±24.6	±28.0	±31.8	±31.2	±35.4	±36.1	±37.1

*: P<0.05, **: P<0.01 (significantly different from control).

Table 3 - continued

Body weights
Male, Female

Study No. P030097

Sex	Group and dose		Body weight (g) on day				
			260	288	316	344	364
Male	Control	N	10	10	10	10	10
		Mean	756.3	786.6	807.1	832.0	843.4
		S.D.	±117.4	±131.2	±136.9	±145.8	±152.9
	0.1 mg/kg	N	9	9	9	9	8
		Mean	748.2	768.4	786.2	795.8	811.9
		S.D.	±90.8	±101.6	±114.1	±129.1	±144.5
	0.5 mg/kg	N	10	10	10	10	10
		Mean	790.8	813.2	832.0	858.3	862.3
		S.D.	±112.4	±118.0	±124.8	±136.1	±139.3
	2.5 mg/kg	N	10	10	10	10	10
		Mean	609.5**	620.7**	623.8**	643.3**	639.4**
		S.D.	±84.2	±88.2	±85.5	±97.1	±96.7
Female	Control	N	10	10	10	10	10
		Mean	394.5	404.6	417.1	430.5	440.1
		S.D.	±66.0	±71.8	±79.9	±85.6	±87.7
	0.5 mg/kg	N	10	10	10	10	10
		Mean	396.2	414.7	433.8	448.5	457.9
		S.D.	±53.5	±57.5	±63.3	±70.3	±73.4
	2.5 mg/kg	N	10	10	10	10	10
		Mean	432.8	446.8	471.1	484.6	495.0
		S.D.	±77.2	±83.2	±98.0	±99.8	±105.5
	12.5 mg/kg	N	10	10	10	10	9
		Mean	390.3	402.3	416.8	426.2	442.4
		S.D.	±41.4	±48.9	±54.3	±69.0	±72.2

**: P<0.01 (significantly different from control).

Table 4 Food consumption
Male, Female

Study No. PQ30097

Sex	Group and dose	Food consumption(g) on day										
		2	8	15	22	29	36	43	50	57 _g	64	
Male	Control	N	20	20	20	20	20	20	20	20	20	20
		Mean	23.1	23.7	24.6	23.3	23.1	23.1	23.4	22.8	22.6	21.5
		S.D.	±2.1	±2.2	±2.9	±3.1	±3.1	±2.3	±2.6	±2.4	±3.2	±3.6
	0.1 mg/kg	N	20	20	20	20	20	20	20	20	20	20
		Mean	22.7	25.3	24.4	24.5	23.4	22.7	22.5	23.2	22.5	21.1
		S.D.	±1.8	±2.4	±2.6	±2.6	±2.4	±2.3	±2.6	±2.7	±2.1	±2.6
	0.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20
		Mean	23.5	25.8*	25.8	23.6	24.3	24.2	24.3	24.3	24.2	23.7*
		S.D.	±1.9	±2.1	±2.8	±2.5	±2.1	±1.8	±1.9	±2.4	±2.2	±2.2
	2.5 mg/kg	N	20	20	20	20	20	20	20	20	19	19
		Mean	23.0	23.7	23.2	22.4	21.7	23.0	21.4	22.5	22.5	21.7
		S.D.	±2.3	±2.1	±3.1	±2.9	±2.8	±3.1	±3.6	±3.8	±3.7	±3.0
Female	Control	N	20	20	20	20	20	20	20	20	20	20
		Mean	16.5	16.5	16.3	15.9	15.9	16.1	15.7	14.2	15.0	13.4
		S.D.	±2.3	±2.8	±2.1	±2.0	±3.0	±3.2	±2.6	±2.4	±2.2	±2.6
	0.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20
		Mean	16.6	16.3	15.5	15.2	15.2	15.4	15.0	15.2	14.9	12.6
		S.D.	±1.7	±2.8	±2.2	±1.5	±1.9	±3.1	±2.5	±2.4	±2.4	±2.6
	2.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20
		Mean	16.4	16.7	15.5	15.4	16.1	16.2	14.8	16.0	14.6	12.9
		S.D.	±1.7	±1.8	±2.3	±1.9	±1.9	±2.0	±3.2	±2.2	±1.8	±2.4
	12.5 mg/kg	N	20	20	20	20	20	20	20	20	20	20
		Mean	16.5	17.1	16.3	15.4	15.8	16.0	14.8	16.0	14.9	14.2
		S.D.	±1.8	±2.1	±2.2	±2.4	±2.9	±2.6	±1.8	±2.7	±2.3	±3.0

*: P<0.05 (significantly different from control).

Table 4 - continued Food consumption
Male, Female.

Study No. P030097

Sex	Group and dose	Food consumption (g) on day										
		71	78	85	91	92	120	148	176	204	232	
Male	Control	N	20	20	20	20	10	10	10	10	10*	10
		Mean	20.0	21.0	20.9	20.7	20.9	20.8	20.7	21.7	20.1	16.5
		S.D.	±2.6	±3.2	±4.0	±3.1	±3.9	±3.4	±4.3	±4.0	±3.0	±2.1
	0.1 mg/kg	N	20	20	20	20	10	10	10	10	10	9
		Mean	19.4	21.5	20.6	19.9	19.6	19.1	20.6	19.8	18.9	17.3
		S.D.	±2.7	±2.4	±2.3	±2.9	±3.2	±4.6	±3.0	±2.2	±3.4	±2.3
	0.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	22.5*	23.8**	22.1	22.2	21.1	22.0	23.7	23.0	22.9	19.4*
		S.D.	±2.3	±2.4	±2.8	±2.1	±5.9	±2.1	±2.5	±1.9	±1.9	±1.7
	2.5 mg/kg	N	19	19	19	19	10	10	10	10	10	10
		Mean	21.4	22.4	21.8	21.3	23.4	25.1*	24.5	24.3	23.8*	21.5**
		S.D.	±2.9	±3.1	±3.2	±2.1	±2.7	±3.3	±5.7	±4.4	±3.4	±3.3
Female	Control	N	20	20	20	20	10	10	10	10	10	10
		Mean	14.1	14.6	14.4	15.5	13.4	13.2	14.0	15.2	13.5	12.2
		S.D.	±2.8	±3.7	±2.9	±2.5	±2.8	±2.3	±2.8	±3.7	±3.0	±3.2
	0.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	13.3	15.2	14.0	13.7	13.9	13.8	15.0	14.6	13.8	12.0
		S.D.	±2.5	±2.0	±2.0	±2.9	±1.6	±2.0	±2.3	±1.7	±2.1	±2.6
	2.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	12.5	15.4	14.3	14.5	14.4	14.2	14.7	15.1	13.5	12.1
		S.D.	±1.8	±1.9	±2.1	±2.9	±2.0	±1.9	±1.7	±3.0	±2.5	±2.4
	12.5 mg/kg	N	20	20	20	20	10	10	10	10	10	10
		Mean	13.6	15.5	14.1	14.9	13.8	13.9	16.1	15.4	15.0	12.5
		S.D.	±2.2	±2.2	±2.5	±2.5	±2.3	±3.2	±2.6	±2.2	±2.4	±2.4

*: P<0.05, **: P<0.01 (significantly different from control).

Table 4 - continued Food consumption
Male, Female

Study No. P030097

Sex	Group and dose	Food consumption (g) on day					
		260	288	316	344	364	
Male	Control	N	10	10	10	10	10
		Mean	17.4	17.5	18.5	19.4	14.9
		S.D.	±2.0	±3.5	±3.3	±5.0	±3.9
	0.1 mg/kg	N	9	9	9	9	8
		Mean	17.3	16.1	15.9	14.1*	13.8
		S.D.	±3.3	±2.2	±2.6	±4.5	±4.3
	0.5 mg/kg	N	10	10	10	10	10
		Mean	19.9	20.1	19.0	20.5	17.7
		S.D.	±2.8	±2.2	±3.4	±3.6	±2.1
	2.5 mg/kg	N	10	10	10	10	10
		Mean	22.0**	23.1**	21.5	22.5	20.8**
		S.D.	±2.1	±3.3	±3.5	±3.9	±2.5
Female	Control	N	10	10	10	10	10
		Mean	13.3	12.7	12.4	13.9	12.0
		S.D.	±3.5	±3.1	±2.7	±1.8	±1.9
	0.5 mg/kg	N	10	10	10	10	10
		Mean	12.8	13.9	14.7	14.0	13.4
		S.D.	±2.1	±2.5	±1.6	±2.3	±1.1
	2.5 mg/kg	N	10	10	10	10	10
		Mean	13.8	11.5	14.5	15.2	13.2
		S.D.	±2.7	±1.4	±2.1	±2.2	±1.5
	12.5 mg/kg	N	10	10	10	10	9
		Mean	14.8	12.4	14.2	15.2	14.0*
		S.D.	±2.8	±1.8	±2.9	±5.0	±2.0

*: P<0.05, **: P<0.01 (significantly different from control).

105
74

Table 5. Urinary findings
Male, Female, 13w

Sex	Group and dose		Urine volume	Osmotic pressure	Specific gravity	Na	K	Cl
			(mL/24hr)	(Osm/kg)		(mEq/24hr)	(mEq/24hr)	(mEq/24hr)
Male	Control	N	10	10	10	10	10	10
		Mean	14.1	1.468	1.050	0.990	2.103	1.302
		S.D.	±8.7	±0.661	±0.022	±0.357	±0.295	±0.380
	0.1 mg/kg	N	10	10	10	10	10	10
		Mean	15.6	1.157	1.040	1.042	2.123	1.444
		S.D.	±5.5	±0.362	±0.012	±0.372	±0.531	±0.440
	0.5 mg/kg	N	10	10	10	10	10	10
		Mean	10.6	1.861	1.062	0.984	2.134	1.320
		S.D.	±2.9	±0.417	±0.014	±0.236	±0.282	±0.198
	2.5 mg/kg	N	9	9	9	9	9	9
		Mean	7.8	2.197**	1.072**	1.147	1.909	1.268
		S.D.	±1.8	±0.382	±0.011	±0.296	±0.372	±0.300
Female	Control	N	10	10	10	10	10	10
		Mean	8.5	1.602	1.051	0.782	1.659	1.143
		S.D.	±5.2	±0.637	±0.019	±0.323	±0.761	±0.470
	0.5 mg/kg	N	10	10	10	10	10	10
		Mean	8.7	1.234	1.040	0.599	1.390	0.878
		S.D.	±2.7	±0.219	±0.008	±0.250	±0.488	±0.345
	2.5 mg/kg	N	10	10	10	10	10	10
		Mean	10.0	1.185	1.039	0.661	1.607	0.973
		S.D.	±3.3	±0.285	±0.009	±0.296	±0.495	±0.285
	12.5 mg/kg	N	10	10	10	10	10	10
		Mean	7.8	1.558	1.051	0.655	1.506	0.952
		S.D.	±2.8	±0.506	±0.016	±0.195	±0.444	±0.305

** : P<0.01 (significantly different from control).

Table 5 - continued

Urinary findings,
Male, Female, 13w

Study No. P030097

Sex	Group and dose	Number of animals	Color		pH							Protein				Glucose
			PY	Y	5.5	6.0	6.5	7.0	7.5	8.0	8.5	-	±	+	++	-
Male	Control	10	3	7	0	0	0	1	2	6	1	1	5	4	0	10
	0.1 mg/kg	10	3	7	1	0	0	0	0	6	3	1	4	5	0	10
	0.5 mg/kg	10	0	10	0	0	0	0	1	9	0	2	2	5	1	10
	2.5 mg/kg	9	0	9	0	0	0	0	1	5	3	1	4	2	2	9
Female	Control	10	0	10	0	1	1	1	1	5	1	8	0	2	0	10
	0.5 mg/kg	10	0	10	0	0	0	1	1	2	6	10	0	0	0	10
	2.5 mg/kg	10	0	10	0	2	0	0	2	3	3	8	2	0	0	10
	12.5 mg/kg	10	0	10	0	0	0	0	3	5	2	7	3	0	0	10

Not significantly different from control.

Abbreviation: PY, pale yellow; Y, yellow.

Grade sign: -, none; ±, trace; +, slight; ++, moderate; +++, severe; +++++, very severe.