

Table 13 Sperm number and motility in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Number of testis sperms		Number of epididymal sperms		% Motile	% Pro- gressive	Swimming speed			Swimming pattern				
			$10^6/\text{testis}$	$10^6/\text{g testis}$	$10^6/\text{cauda}$	$10^6/\text{g cauda}$			VAP	VSL	VCL	ALH	BCF	STR	LIN	
F0	Control	24	Mean	184.1	111.4	268.5	856.4	88.1	70.9	159.6	112.1	365.7	20.1	27.9	69.3	30.4
			S.D.	29.3	13.2	47.6	94.4	9.3	17.4	20.8	22.5	53.4	1.1	1.5	6.6	2.8
	DCBS 80 ppm	23	Mean	187.7	110.7	276.2	838.9	92.6	77.3	159.8	114.1	370.1	19.9	27.4	70.7	30.7
			S.D.	28.3	15.7	40.3	99.4	8.2	15.3	19.2	20.0	42.5	1.1	1.5	5.7	3.0
	DCBS 600 ppm	24	Mean	184.2	110.6	269.9	850.3	93.2	77.4	162.7	116.1	372.3	20.0	27.6	71.0	31.3
			S.D.	32.7	17.1	56.8	122.1	5.9	12.1	22.0	19.3	49.8	1.3	2.2	4.3	2.5
	DCBS 4500 ppm	24	Mean	180.8	106.1	263.7	844.2	89.4	70.5	156.8	110.5	358.4	19.9	28.3	69.5	30.6
			S.D.	35.4	18.8	62.8	191.3	10.2	22.2	25.3	29.2	56.3	1.0	2.3	8.6	4.0
F1	Control	24	Mean	194.5	115.3	273.6	849.9	92.3	81.8	175.2	126.9	399.5	21.3	26.4	72.5	32.0
			S.D.	23.0	9.5	40.0	69.4	5.0	8.1	9.8	10.2	19.8	0.9	1.6	3.3	2.1
	DCBS 80 ppm	24	Mean	181.1	108.4	254.0	821.5	92.9	81.8	171.7	123.9	391.5	20.9	26.8	72.1	31.9
			S.D.	21.3	14.3	40.4	106.8	4.0	4.9	11.2	10.3	28.6	0.8	1.4	2.7	2.0
	DCBS 600 ppm	24	Mean	186.3	111.1	256.2	827.2	93.3	83.9	172.4	126.0	395.1	20.8	26.1	73.3	32.1
			S.D.	22.5	11.3	46.0	93.3	5.6	6.4	11.4	10.5	28.6	0.8	1.6	2.9	1.8
	DCBS 4500 ppm	24	Mean	201.0	113.6	250.3	807.0	93.0	82.7	171.3	125.7	393.6	20.5 *	27.0	73.5	32.2
			S.D.	33.3	15.0	55.4	127.5	7.4	8.2	13.9	12.6	29.8	1.0	1.8	2.8	1.5

VAP: Mean path velocity ($\mu\text{m/sec}$).

VSL: Straight line average velocity ($\mu\text{m/sec}$).

VCL: Mean curvilinear velocity ($\mu\text{m/sec}$).

ALH: Mean lateral head displacement (μm).

BCF: Mean beat cross frequency (Hz).

STR: Mean straightness (%) = VSL/VAP x 100.

LIN: Mean linearity (%) = VSL/VCL x 100.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

Table 14 Abnormal sperm ratio in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Total	Abnormal sperm ratio (%)											
				Head						Neck and middle piece				Tail	
				Tailless sperm	Small sized	Banana shaped	Hooked shaped	Truncated shaped	Amorphous shaped	Two heads	Flexion	Two necks	Enlarge- ment	Fragmen- tation	Two tails
F0	Control	24	Mean	1.1	1.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	0.6	0.6	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	DCBS 80 ppm	23	Mean	1.2	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	0.8	0.8	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
	DCBS 600 ppm	24	Mean	2.4	2.2	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
			S.D.	3.5	3.5	0.0	0.2	0.0	0.1	0.1	0.0	0.2	0.0	0.0	0.0
	DCBS 4500 ppm	24	Mean	2.0	1.8	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	2.4	2.0	0.0	0.5	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
F1	Control	24	Mean	1.4	1.3	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	1.3	1.2	0.0	0.3	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0
	DCBS 80 ppm	24	Mean	1.1	0.9	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	0.8	0.8	0.0	0.2	0.0	0.2	0.1	0.0	0.1	0.0	0.0	0.0
F1	DCBS 600 ppm	24	Mean	1.2	1.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	1.7	1.6	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.0
F1	DCBS 4500 ppm	24	Mean	1.6	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			S.D.	1.9	1.8	0.1	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0

46
78

Table 15 Sexual development in F1 parental rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Male				Female			
		Number of animals	Age at preputial separation (days)	Body weight (g) on the day at preputial separation	Number of animals	Age at vaginal opening (days)	Body weight (g) on the day at vaginal opening		
F1	Control	24	Mean S.D.	41.3 1.6	226.9 20.3	24	Mean S.D.	29.6 1.0	104.6 9.4
	DCBS 80 ppm	24	Mean S.D.	41.4 1.6	226.5 18.5	24	Mean S.D.	30.0 1.7	109.1 10.6
	DCBS 600 ppm	24	Mean S.D.	41.8 1.6	228.3 17.0	24	Mean S.D.	31.2 ^{ss} 1.7	112.1 * 13.8
	DCBS 4500 ppm	24	Mean S.D.	42.8 ** 1.5	229.6 17.5	24	Mean S.D.	31.1 ^{ss} 1.3	112.3 * 9.1

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

^{ss}: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

Table 16 Locomotor activity count in F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Locomotor activity count							
			Determination time (minutes)							
			0-10	10-20	20-30	30-40	40-50	50-60	0-60	
F1	Control	10	Mean	206.0	76.8	26.3	14.3	5.3	0.8	329.5
			S.D.	58.9	40.7	25.0	30.6	16.8	2.2	126.7
	DCBS 80 ppm	10	Mean	248.8	63.1	34.2	9.6	11.0	1.6	368.3
			S.D.	78.8	35.7	32.6	14.3	19.7	5.1	146.2
	DCBS 600 ppm	10	Mean	208.7	78.9	58.1	52.2	12.0	14.7	424.6
			S.D.	111.6	63.7	71.3	89.1	28.2	31.6	230.1
	DCBS 4500 ppm	10	Mean	204.6	72.2	20.9	11.5	1.0	0.0	310.2
			S.D.	154.5	86.4	20.1	17.0	2.5	0.0	236.3

Table 17 Locomotor activity count in F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Locomotor activity count							
			Determination time (minutes)							
			0-10	10-20	20-30	30-40	40-50	50-60	0-60	
F1	Control	10	Mean	168.3	55.7	23.6	10.1	1.8	0.5	260.0
			S.D.	66.1	42.3	27.4	20.6	5.7	1.6	115.6
	DCBS 80 ppm	10	Mean	212.0	59.3	34.8	19.7	11.2	0.0	337.0
			S.D.	121.3	46.5	75.6	54.4	35.4	0.0	297.9
	DCBS 600 ppm	10	Mean	220.2	74.3	25.7	9.1	4.5	0.8	334.6
			S.D.	103.7	48.9	32.7	26.1	11.7	2.5	181.6
	DCBS 4500 ppm	10	Mean	203.1	52.6	19.7	14.5	6.6	0.0	296.5
			S.D.	109.8	42.2	21.8	34.9	20.9	0.0	173.4

Table 18 Data on learning tests in F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Time for maze trials (sec)				Number of errors for maze trials				
			Day 1	Day 2	Day 3	Day 4	Day 1	Day 2	Day 3	Day 4	
F1	Control	10	Mean	8.2	58.0	29.0	30.9	0.0	10.2	2.8	2.8
			S.D.	2.4	35.1	9.8	12.1	0.1	7.2	1.8	2.0
	DCBS 80 ppm	10	Mean	8.1	47.1	23.6	19.8	0.1	8.5	3.0	1.3
			S.D.	1.2	20.5	7.0	7.9	0.1	2.2	1.7	0.9
	DCBS 600 ppm	10	Mean	8.9	46.3	35.1	26.7	0.1	7.8	5.1	2.6
			S.D.	1.9	19.6	16.1	11.8	0.1	2.9	3.9	2.2
	DCBS 4500 ppm	10	Mean	8.7	58.2	30.7	27.0	0.0	10.8	4.3	2.4
			S.D.	2.1	23.1	11.5	7.6	0.1	4.1	2.5	1.7

Day 1 : Used a straight channel.

Days 2-4 : Used a multiple T-maze.

Table 19 Data on learning tests in F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Time for maze trials (sec)				Number of errors for maze trials				
			Day 1	Day 2	Day 3	Day 4	Day 1	Day 2	Day 3	Day 4	
F1	Control	10	Mean	10.9	36.8	31.8	19.8	0.2	6.1	4.5	2.9
			S.D.	3.4	12.4	21.7	5.8	0.2	2.6	3.3	1.4
	DCBS 80 ppm	10	Mean	8.8	41.9	27.6	25.2	0.1	6.8	3.9	3.6
			S.D.	2.6	15.1	12.9	13.1	0.2	2.5	2.5	4.3
	DCBS 600 ppm	10	Mean	9.7	57.7 *	36.6	28.1	0.2	8.3	5.3	3.7
			S.D.	4.0	18.7	17.9	14.2	0.4	2.7	2.9	3.4
	DCBS 4500 ppm	10	Mean	10.2	57.7 *	39.5	31.5	0.2	9.4 *	5.6	3.2
			S.D.	2.5	17.8	17.7	18.7	0.2	2.8	3.4	2.8

Day 1 : Used a straight channel.

Days 2-4 : Used a multiple T-maze.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

Table 20 Hematological findings in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	WBC $10^3/\mu\text{L}$	Differential count of WBC %						
				Neutrophil		Eosino- phil	Baso- phil	Mono- cyte	Lympho- cyte	Others
				Stab form	Seg- mented					
F0	Control	10	Mean	92	0.96	12.64	1.16	0.00	2.72	82.5
			S.D.	22	0.63	5.11	0.64	0.00	1.08	5.2
	DCBS 80 ppm	10	Mean	83	0.92	8.88	1.12	0.00	2.52	86.6
			S.D.	17	0.33	2.38	0.75	0.00	0.84	2.4
	DCBS 600 ppm	10	Mean	96	0.88	9.28	1.84	0.00	2.48	85.5
			S.D.	26	0.41	3.62	0.97	0.00	0.92	3.5
	DCBS 4500 ppm	10	Mean	111	0.52	9.12	1.44	0.00	2.00	86.9 *
			S.D.	32	0.53	3.28	0.85	0.00	0.68	3.9
F1	Control	10	Mean	104	0.76	11.16	1.24	0.00	2.64	84.2
			S.D.	17	0.40	3.45	0.87	0.00	0.95	4.5
	DCBS 80 ppm	10	Mean	103	0.68	9.96	1.12	0.00	3.28	85.0
			S.D.	13	0.50	3.71	0.73	0.00	1.46	4.9
	DCBS 600 ppm	10	Mean	126	0.84	9.32	0.92	0.08	2.44	86.4
			S.D.	33	0.48	4.68	0.60	0.17	1.43	6.1
	DCBS 4500 ppm	10	Mean	109	0.48	8.96	0.76	0.04	2.64	87.1
			S.D.	25	0.37	3.91	0.61	0.13	0.87	3.5

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

Table 21 Hematological findings in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	WBC $10^3/\mu\text{L}$	Differential count of WBC %						
				Neutrophil		Eosino- phil	Baso- phil	Mono- cyte	Lympho- cyte	Others
				Stab form	Seg- mented					
F0	Control	10	Mean	87	1.56	20.68	2.20	0.04	4.44	71.1
			S.D.	19	0.91	6.78	1.44	0.13	1.42	7.8
	DCBS 80 ppm	10	Mean	74	1.72	21.40	1.44	0.00	3.48	72.0
			S.D.	17	0.92	5.71	0.95	0.00	1.46	6.2
	DCBS 600 ppm	10	Mean	98	1.24	22.08	1.24	0.00	3.44	72.0
			S.D.	21	0.74	8.59	0.61	0.00	2.16	10.1
	DCBS 4500 ppm	10	Mean	76	1.68	21.92	1.60	0.00	3.52	71.3
			S.D.	17	0.96	10.81	0.82	0.00	1.63	11.8
F1	Control	10	Mean	98	1.48	19.52	1.00	0.04	3.40	74.6
			S.D.	20	0.50	5.76	0.43	0.13	2.02	5.9
	DCBS 80 ppm	10	Mean	90	1.12	19.72	0.72	0.00	2.60	75.8
			S.D.	17	0.65	8.13	0.59	0.00	1.10	8.2
	DCBS 600 ppm	10	Mean	96	1.28	13.48	0.48	0.04	1.88	82.8 *
			S.D.	26	0.73	6.32	0.56	0.13	1.07	6.7
	DCBS 4500 ppm	10	Mean	98	1.24	17.40	0.84	0.00	3.04	77.5
			S.D.	24	0.58	4.58	0.61	0.00	2.18	6.0

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

Table 22 Blood chemical findings in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals		TP g/dL	Albumin g/dL	Globulin g/dL
F0	Control	10	Mean	6.27	2.43	3.84
			S.D.	0.32	0.13	0.31
	DCBS 80 ppm	10	Mean	6.31	2.43	3.88
			S.D.	0.21	0.09	0.18
	DCBS 600 ppm	10	Mean	6.33	2.40	3.93
			S.D.	0.31	0.12	0.28
	DCBS 4500 ppm	10	Mean	6.54	2.49	4.05
			S.D.	0.28	0.07	0.24
F1	Control	10	Mean	6.07	2.29	3.78
			S.D.	0.26	0.12	0.24
	DCBS 80 ppm	10	Mean	6.15	2.31	3.84
			S.D.	0.31	0.11	0.25
	DCBS 600 ppm	10	Mean	6.26	2.31	3.95
			S.D.	0.23	0.13	0.29
	DCBS 4500 ppm	10	Mean	6.16	2.31	3.85
			S.D.	0.26	0.07	0.27

Table 23 Blood chemical findings in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals		TP g/dL	Albumin g/dL	Globulin g/dL
F0	Control	10	Mean	6.45	2.71	3.74
			S.D.	0.27	0.14	0.25
	DCBS 80 ppm	10	Mean	6.80	2.78	4.02
			S.D.	0.43	0.19	0.29
	DCBS 600 ppm	10	Mean	6.28	2.60	3.68
			S.D.	0.30	0.17	0.22
	DCBS 4500 ppm	10	Mean	6.45	2.65	3.80
			S.D.	0.39	0.16	0.25
F1	Control	10	Mean	6.36	2.66	3.70
			S.D.	0.33	0.13	0.25
	DCBS 80 ppm	10	Mean	6.34	2.61	3.73
			S.D.	0.41	0.14	0.27
	DCBS 600 ppm	10	Mean	6.23	2.59	3.64
			S.D.	0.37	0.20	0.25
	DCBS 4500 ppm	10	Mean	6.39	2.63	3.76
			S.D.	0.33	0.15	0.23

55
187

Table 24 Serum hormone levels in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals		Testosterone (ng/mL)	DHT ^a (pg/mL)	LH ^b (ng/mL)	FSH (ng/mL)
F0	Control	8	Mean	1.32	80.9	1.03	8.87
			S.D.	0.90	41.4	0.10	1.61
	DCBS 80 ppm	8	Mean	1.38	82.2	0.83	9.55
			S.D.	0.97	32.0	0.72	2.16
F1	DCBS 600 ppm	8	Mean	1.01	69.1	1.26	9.43
			S.D.	0.29	28.3	0.38	1.40
	DCBS 4500 ppm	8	Mean	1.07	56.4	1.08	9.31
			S.D.	0.92	34.9	0.54	0.97
F1	Control	8	Mean	0.40	39.2	1.38	7.68
			S.D.	0.13	26.7	0.32	1.46
	DCBS 80 ppm	8	Mean	0.99 ^s	49.7	1.23	7.05
			S.D.	0.70	14.6	0.56	1.02
F1	DCBS 600 ppm	8	Mean	0.89	48.1	1.72 ^s	6.92
			S.D.	0.63	21.1	0.12	1.41
	DCBS 4500 ppm	8	Mean	0.72	41.9	1.58	7.84
			S.D.	0.43	25.6	0.34	1.53

a: The actual measurement of DHT was below the lower limit of quantification (<25.0 pg/mL) in one F1 animal each in the control and 4500 ppm groups.

b: The actual measurement of LH was below the lower limit of quantification (<0.80 ng/mL) in three F0 and one F1 animals in the 80 ppm group and one F0 animal in the 4500 ppm group.

^s: Significantly different from the control at p≤0.05 by Mann-Whitney U-test.

Table 25 Serum hormone levels in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals		Estradiol (pg/mL)	Progesterone (ng/mL)	LH ^a (ng/mL)	FSH (ng/mL)
F0	Control	8	Mean	80.5	5.17	1.30	5.75
			S.D.	15.7	3.59	0.22	0.99
	DCBS 80 ppm	8	Mean	66.7	8.49	1.13	6.29
			S.D.	16.2	4.37	0.27	1.14
F1	DCBS 600 ppm	8	Mean	68.7	8.08	1.35	5.99
			S.D.	16.0	4.70	0.20	1.06
	DCBS 4500 ppm	8	Mean	64.0	9.88	1.15	5.60
			S.D.	12.3	4.20	0.49	0.86
F1	Control	8	Mean	73.8	6.42	1.45	3.12
			S.D.	25.3	2.84	0.26	0.28
	DCBS 80 ppm	8	Mean	61.9	8.95	1.34	3.63
			S.D.	17.2	3.55	0.59	1.38
F1	DCBS 600 ppm	8	Mean	59.7	7.20	1.52	3.91
			S.D.	21.5	4.66	0.24	1.32
	DCBS 4500 ppm	8	Mean	63.5	8.59	1.55	4.50
			S.D.	26.7	5.85	0.42	1.50

a: The actual measurement of LH was below the lower limit of quantification (<0.80 ng/mL) in one F1 animal in the 80 ppm group and one F0 animal in the 4500 ppm group.

Table 26 Autopsy findings in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Item	Control				DCBS (ppm)											
						80				600				4500			
		A	B	C	T	A	B	C	T	A	B	C	T	A	B	C	T
F0	Number of animals examined	22	2	0	24	23	0	1	24	24	0	0	24	24	0	0	24
	Number of animals with abnormal findings	0	0	-	0	0	-	1	1	3	-	-	3	1	-	-	1
	Findings ^a																
	Incisor: Malocclusion	0	0	-	0	0	-	1	1	1	-	-	1	0	-	-	0
	Nasal bone: Fracture	0	0	-	0	0	-	1	1	0	-	-	0	0	-	-	0
	Stomach, duodenum, jejunum, ileum and cecum: Retention gas	0	0	-	0	0	-	1	1	0	-	-	0	0	-	-	0
	Ileum: White mass, serosa	0	0	-	0	0	-	0	0	1	-	-	1	0	-	-	0
	Thickening, mucosa	0	0	-	0	0	-	0	0	1	-	-	1	0	-	-	0
	Mesenteric lymph node: Swelling	0	0	-	0	0	-	0	0	1	-	-	1	0	-	-	0
	Testis: Atrophy	0	0	-	0	0	-	0	0	0	-	-	0	1	-	-	1
	Epididymis: Atrophy	0	0	-	0	0	-	0	0	0	-	-	0	1	-	-	1
F1	Number of animals examined	23	1	0	24	22	2	0	24	20	4	0	24	24	0	0	24
	Number of animals with abnormal findings	2	1	-	3	4	0	-	4	1	0	-	1	6	-	-	6
	Findings ^a																
	Incisor: Malocclusion	0	0	-	0	0	0	-	0	0	0	-	0	1	-	-	1
	Ileum: Diverticulum	0	0	-	0	0	0	-	0	0	0	-	0	2	-	-	2
	Kidney : Dilatation, renal pelvis	1	1	-	2	3	0	-	3	1	0	-	1	2	-	-	2
	Fine white granule, renal pelvis	0	0	-	0	1	0	-	1	0	0	-	0	0	-	-	0
	Seminal vesicle : Small size	0	0	-	0	1	0	-	1	0	0	-	0	1	-	-	1
	Testis: Atrophy	1	0	-	1	0	0	-	0	0	0	-	0	0	-	-	0
	Epididymis: Atrophy	1	0	-	1	0	0	-	0	0	0	-	0	0	-	-	0

Statistical analyses were made based on the total number of animals examined.

Fate: A, animals that impregnated a female; B, animals that unsuccessfully mated or did not impregnate a female; C, animals that were euthanized during the study; T, total (A+B+C).

a: Values represent the number of animals that showed abnormal findings.

-: Not applicable.

5
8
0

Table 27 Autopsy findings in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Item	Control				DCBS (ppm)											
						80				600				4500			
		A	B	C	T	A	B	C	T	A	B	C	T	A	B	C	T
F0	Number of animals examined	22	2	0	24	24	0	0	24	24	0	0	24	24	0	0	24
	Number of animals with abnormal findings	4	2	-	6	0	-	-	0 *	0	-	-	0 *	1	-	-	1
	Findings ^a																
	Skin: Subcutaneous yellowish/grayish white mass	1	0	-	1	0	-	-	0	0	-	-	0	1	-	-	1
	Incisor: Malocclusion/fracture	3	0	-	3	0	-	-	0	0	-	-	0	0	-	-	0
	Nasal bone: Deformity	1	0	-	1	0	-	-	0	0	-	-	0	0	-	-	0
	Ileum: Diverticulum	0	1	-	1	0	-	-	0	0	-	-	0	0	-	-	0
	Uterine horn: Retention, yellowish white mucous fluid	0	1	-	1	0	-	-	0	0	-	-	0	0	-	-	0
	Vagina: Atresia	0	2	-	2	0	-	-	0	0	-	-	0	0	-	-	0
F1	Number of animals examined	22	1	1	24	22	2	0	24	21	3	0	24	23	1	0	24
	Number of animals with abnormal findings	0	0	0	0	0	0	-	0	1	0	-	1	4	1	-	5 *
	Findings ^a																
	Incisor: Malocclusion	0	0	0	0	0	0	-	0	0	0	-	0	3	0	-	3
	Cerebrum: Dilatation, ventricle	0	0	0	0	0	0	-	0	0	0	-	0	0	1	-	1
	Thyroid: Aplasia	0	0	0	0	0	0	-	0	0	0	-	0	1	0	-	1
	Thymus: Atrophy	0	0	0	0	0	0	-	0	0	0	-	0	1	0	-	1
	Kidney: Dilatation, renal pelvis	0	0	0	0	0	0	-	0	1	0	-	1	0	0	-	0
	Ureter: Dilatation	0	0	0	0	0	0	-	0	1	0	-	1	0	0	-	0

Statistical analyses were made based on the total number of animals examined.

Fate: A, animals that had weanlings; B, animals that were not pregnant and that did not produce viable pups or weanlings; C, animals that died during the study; T, total (A+B+C).

a: Values represent the number of animals that showed abnormal findings.

-: Not applicable.

*: Significantly different from the control at $p \leq 0.05$ by Fisher's exact probability test.

Table 28 Absolute and relative organ weights in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Body weight		Brain		Pituitary gland		Thyroid ^a		Thymus		Liver		Kidney ^a		Spleen	
			g	g	g	%	mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	g	%	g	%	mg	10 ⁻³ %
F0	Control	24	Mean	624.4	2.25	0.366	12.8	2.07	24.0	3.85	300	48.2	20.45	3.25	3.64	0.585	882	142
			S.D.	76.0	0.08	0.041	1.3	0.22	4.0	0.60	73	11.3	4.34	0.33	0.42	0.043	158	20
	DCBS 80 ppm	23	Mean	612.5	2.28	0.375	13.1	2.16	23.7	3.87	289	47.2	19.76	3.22	3.64	0.595	835	137
			S.D.	61.0	0.10	0.034	1.4	0.22	4.3	0.59	55	8.2	2.94	0.25	0.41	0.044	143	20
	DCBS 600 ppm	24	Mean	603.4	2.25	0.376	12.9	2.15	24.4	4.08	318	52.9	19.76	3.27	3.57	0.593	846	140
			S.D.	51.2	0.07	0.031	1.3	0.21	4.1	0.81	67	11.1	2.09	0.18	0.29	0.035	167	24
	DCBS 4500 ppm	24	Mean	575.4 ^s	2.25	0.392 *	12.4	2.15	26.6	4.63 **	272	47.2	19.96	3.47 ^s	3.69	0.641 **	748 **	131
			S.D.	42.3	0.08	0.030	1.2	0.19	5.2	0.86	49	6.8	2.21	0.26	0.36	0.040	112	21
F1	Control	24	Mean	630.7	2.26	0.363	13.6	2.17	24.9	3.95	346	54.8	20.80	3.28	3.70	0.586	909	145
			S.D.	74.7	0.10	0.038	1.4	0.23	4.9	0.66	116	17.0	3.73	0.29	0.52	0.041	129	16
	DCBS 80 ppm	24	Mean	605.1	2.29	0.380	13.9	2.30	23.3	3.86	269 ^s	44.5 ^s	19.69	3.25	3.66	0.606	845	139
			S.D.	47.7	0.06	0.028	1.3	0.25	4.7	0.78	54	8.9	2.32	0.19	0.23	0.042	141	18
DCBS 600 ppm	24	Mean	614.2	2.26	0.370	13.9	2.27	23.8	3.88	331	53.9	21.19	3.46	3.69	0.602	847	138	
			S.D.	52.5	0.06	0.030	1.1	0.17	4.5	0.69	83	12.7	2.06	0.28	0.36	0.047	124	17
DCBS 4500 ppm	24	Mean	622.6	2.21 ^s	0.356	14.0	2.26	24.6	3.95	316	50.9	22.82 ^s	3.65 **	3.91	0.629 **	869	139	
			S.D.	51.8	0.09	0.027	1.6	0.26	4.9	0.67	62	9.8	3.37	0.28	0.43	0.044	162	17

(to be continued)

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

^s: Significantly different from the control at $p \leq 0.05$ by Mann-Whitney U-test.

Table 28 (continued) Absolute and relative organ weights in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Adrenal ^a		Testis ^a		Epididymis ^a		Seminal vesicle		Prostate		
			mg	10 ⁻³ %	g	%	mg	10 ⁻³ %	g	%	mg	10 ⁻³ %	
F0	Control	24	Mean	58.0	9.3	3.54	0.576	1350	219	2.31	0.376	899	145
			S.D.	9.5	1.3	0.24	0.083	105	26	0.25	0.062	189	32
	DCBS 80 ppm	23	Mean	57.9	9.5	3.62	0.598	1379	228	2.28	0.376	910	149
			S.D.	7.7	1.1	0.20	0.075	74	28	0.24	0.060	150	20
	DCBS 600 ppm	24	Mean	57.7	9.6	3.57	0.594	1354	225	2.27	0.376	853	142
			S.D.	7.4	1.2	0.24	0.048	113	17	0.29	0.041	234	37
	DCBS 4500 ppm	24	Mean	51.0 **	8.9	3.65	0.637 **	1330	232	2.15	0.375	769	134
			S.D.	6.1	1.0	0.31	0.068	115	21	0.21	0.034	165	26
F1	Control	24	Mean	60.5	9.6	3.60	0.575	1348	215	2.30	0.368	838	133
			S.D.	9.8	1.5	0.35	0.062	138	24	0.23	0.047	174	24
	DCBS 80 ppm	24	Mean	60.3	10.0	3.61	0.601	1342	223	2.19	0.364	812	134
			S.D.	7.1	1.0	0.27	0.073	67	21	0.28	0.054	181	28
	DCBS 600 ppm	24	Mean	61.8	10.1	3.60	0.589	1327	217	2.21	0.362	822	134
			S.D.	7.2	1.3	0.27	0.066	111	22	0.22	0.039	190	29
	DCBS 4500 ppm	24	Mean	61.3	9.8	3.78	0.610	1346	217	2.07 **	0.333 *	784	127
			S.D.	13.1	2.0	0.32	0.062	118	19	0.26	0.045	168	31

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

Table 29 Absolute and relative organ weights in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Body weight		Brain		Pituitary gland		Thyroid ^a		Thymus		Liver		Kidney ^a		Spleen	
			g	g	g	%	mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	g	%	g	%	mg	10 ⁻³ %
F0	Control	22	Mean	305.9	2.04	0.670	13.8	4.51	18.8	6.19	237	77.0	13.73	4.48	2.13	0.696	632	207
			S.D.	28.5	0.10	0.050	1.8	0.47	4.4	1.57	71	21.6	2.51	0.67	0.29	0.063	97	27
	DCBS 80 ppm	24	Mean	311.6	2.12 **	0.681	15.2 *	4.87	17.7	5.67	209	67.2	13.83	4.43	2.20	0.707	582	187 *
			S.D.	17.1	0.09	0.046	1.9	0.63	2.2	0.71	55	17.5	1.73	0.44	0.16	0.058	78	26
	DCBS 600 ppm	24	Mean	313.7	2.12 **	0.676	14.7	4.70	16.9	5.40	242	77.7	13.86	4.41	2.29	0.731	583	186 *
			S.D.	17.8	0.07	0.042	2.2	0.66	3.1	0.92	57	19.8	1.84	0.47	0.23	0.068	77	22
	DCBS 4500 ppm	24	Mean	288.5 ^s	2.07	0.720 **	13.6	4.71	17.5	6.10	197	68.5	13.67	4.74	2.27	0.786 **	560 *	194
			S.D.	16.2	0.07	0.045	1.8	0.60	2.2	0.86	59	20.8	1.48	0.44	0.19	0.057	79	27
F1	Control	22	Mean	331.9	2.08	0.632	15.9	4.83	19.0	5.72	251	75.3	14.55	4.39	2.37	0.713	632	191
			S.D.	32.5	0.08	0.056	2.0	0.73	3.9	0.98	69	18.4	1.66	0.28	0.30	0.046	73	18
	DCBS 80 ppm	22	Mean	331.2	2.17 **	0.658	16.1	4.90	18.2	5.51	212	64.1	14.18	4.28	2.39	0.723	599	181
			S.D.	28.5	0.08	0.056	2.4	0.79	2.7	0.70	47	14.2	2.14	0.49	0.22	0.040	63	15
	DCBS 600 ppm	21	Mean	331.3	2.15 *	0.651	15.8	4.78	17.7	5.35	261	79.2	14.32	4.33	2.40	0.726	609	184
			S.D.	23.1	0.08	0.043	1.8	0.52	3.5	1.08	65	20.2	1.49	0.41	0.21	0.063	80	19
	DCBS 4500 ppm	23	Mean	330.2	2.08	0.633	16.1	4.89	19.4	5.89	211	64.0	15.83	4.81 ^{ss}	2.53	0.771 ^{ss}	639	194
			S.D.	30.8	0.08	0.060	1.9	0.66	4.1	1.15	63	18.7	2.11	0.59	0.26	0.080	115	37

(to be continued)

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

^s: Significantly different from the control at $p \leq 0.05$ by Mann-Whitney U-test.

^{ss}: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

62-94-

Table 29 (continued) Absolute and relative organ weights in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Group	Number of animals	Adrenal ^a		Ovary ^a		Uterus		
			mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	
F0	Control	22	Mean	66.8	21.8	100.2	32.8	957	313
			S.D.	9.4	2.3	15.8	4.4	189	56
	DCBS 80 ppm	24	Mean	65.9	21.2	95.4	30.7	962	310
			S.D.	6.1	2.1	11.6	3.9	186	65
	DCBS 600 ppm	24	Mean	68.3	21.8	101.1	32.3	975	311
			S.D.	8.6	2.6	9.1	3.1	198	63
	DCBS 4500 ppm	24	Mean	69.6	24.1 **	91.6	31.8	1024	357
			S.D.	8.1	2.7	12.4	4.1	203	76
F1	Control	22	Mean	70.0	21.2	110.6	33.4	927	280
			S.D.	9.7	3.2	13.0	2.9	191	54
	DCBS 80 ppm	22	Mean	73.5	22.2	109.1	33.0	928	283
			S.D.	10.9	3.1	16.3	4.5	128	48
	DCBS 600 ppm	21	Mean	73.4	22.2	108.5	32.8	976	295
			S.D.	9.3	3.0	12.5	3.2	185	52
	DCBS 4500 ppm	23	Mean	77.5 *	23.6 *	108.2	32.8	949	288
			S.D.	8.9	3.2	13.4	3.3	192	52

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

Table 30 Histopathological findings in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Item	Control			DCBS (ppm)								
		A	B	T	80			600			4500		
F0	Number of animals examined	22	2	24	0	0	0	0	0	0	24	0	24
Findings in organs/tissues examined^{a,b,c}													
Liver: Microgranuloma		4	2	6	-	-	-	-	-	-	5	-	5
Kidney: Hyaline droplet, proximal tubular epithelium		12	2	14	-	-	-	-	-	-	20	-	20
Eosinophilic body, proximal tubular epithelium		12	2	14	-	-	-	-	-	-	20	-	20
Regeneration, tubular epithelium		5	0	5	-	-	-	-	-	-	1	-	1
Cast, hyaline		2	0	2	-	-	-	-	-	-	2	-	2
Cellular infiltration, inflammatory cell, renal pelvic mucosa		0	1	1	-	-	-	-	-	-	0	-	0
Cyst		2	0	2	-	-	-	-	-	-	1	-	1
Testis: Atrophy, seminiferous tubule		0	0	0	-	-	-	-	-	-	1	-	1
Epididymis: Decrease, sperm		0	0	0	-	-	-	-	-	-	1	-	1
Cell debris, lumen		0	0	0	-	-	-	-	-	-	1	-	1
Prostate: Cellular infiltration, inflammatory cell		15	0	15	-	-	-	-	-	-	17	-	17
Pituitary gland: Cyst, pars distalis		1	0	1	-	-	-	-	-	-	0	-	0
Cyst, pars intermedia		1	0	1	-	-	-	-	-	-	0	-	0
Tubular hyperplasia, pars intermedia		1	0	1	-	-	-	-	-	-	1	-	1
Findings in organs/tissues in which abnormal findings were grossly observed^{a,d}													
Ileum: Cyst, squamous cell		-	-	-	-	-	-	1(2)	-	1(2)	-	-	-
Cellular infiltration, inflammatory cell		-	-	-	-	-	-	1(2)	-	1(2)	-	-	-
Mesenteric lymph node: Hyperplasia, lymphoid		-	-	-	-	-	-	1(1)	-	1(1)	-	-	-

(to be continued)

Fate: A, animals that impregnated a female; B, animals that did not impregnate a female; T, total (A+B).

a: Values represent the number of animals that showed abnormal findings.

b: Organs/tissues examined were the liver, kidneys, thymus, thyroids, spleen, adrenals, bone marrow (femur), mesenteric lymph node, Peyer's patch, testis, epididymis, seminal vesicles, coagulating glands, prostate, mammary gland and pituitary gland.

c: Statistical analyses were made between the control and 4500 ppm groups based on the total number of animals examined.

d: Values in parentheses are the number of animals examined.

-: Not applicable.

Table 30 (continued) Histopathological findings in F0 and F1 parental male rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Item	Control			DCBS (ppm)								
					80			600			4500		
		A	B	T	A	B	T	A	B	T	A	B	T
F1	Number of animals examined	23	1	24	0	2	2	0	4	4	24	0	24
Findings in organs/tissues examined^{a,b,c}													
	Liver: Microgranuloma	8	0	8	-	1	1	-	3	3	7	-	7
	Kidney: Hyaline droplet, proximal tubular epithelium	20	0	20	-	1	1	-	2	2	24	-	24
	Eosinophilic body, proximal tubular epithelium	20	0	20	-	1	1	-	2	2	24	-	24
	Regeneration, tubular epithelium	6	1	7	-	0	0	-	1	1	8	-	8
	Cast, hyaline	4	0	4	-	0	0	-	0	0	5	-	5
	Mineralization, papilla	0	0	0	-	0	0	-	0	0	1	-	1
	Dilatation, renal pelvis	1	1	2	-	0	0	-	0	0	2	-	2
	Testis: Atrophy, seminiferous tubule	1	0	1	-	0	0	-	0	0	0	-	0
	Epididymis: Decrease, sperm	1	0	1	-	0	0	-	0	0	0	-	0
	Cell debris, lumen	1	0	1	-	0	0	-	0	0	0	-	0
	Prostate: Cellular infiltration, inflammatory cell	13	0	13	-	2	2	-	1	1	12	-	12
	Pituitary gland: Cyst, pars distalis	0	0	0	-	0	0	-	1	1	1	-	1
	Tubular hyperplasia, pars intermedia	0	0	0	-	0	0	-	0	0	1	-	1

Fate: A, animals that impregnated a female; B, animals that unsuccessfully mated or did not impregnate a female; T, total (A+B).

a: Values represent the number of animals that showed abnormal findings.

b: Organs/tissues examined were the liver, kidneys, thymus, thyroids, spleen, adrenals, bone marrow (femur), mesenteric lymph node, Peyer's patch, testis, epididymis, seminal vesicles, coagulating glands, prostate, mammary gland and pituitary gland.

c: Statistical analyses were made between the control and 4500 ppm groups based on the total number of animals examined.

-: Not applicable.

Table 31 Histopathological findings in F0 and F1 parental female rats treated with N,N-Dicyclohexyl-2-benzothiazolesulfenamide (DCBS) in the two-generation reproductive toxicity study (SR05241)

Gener- ation	Item	Control				DCBS (ppm)							
		A	B	C	T	A	B	C	T	A	B	C	T
F0	Number of animals examined	22	2	0	24	0	0	0	0	0	0	0	24
Findings in organs/tissues examined^{a,b,c}													
	Liver: Microgranuloma	5	0	-	5	-	-	-	-	-	7	-	-
	Kidney: Regeneration, tubular epithelium	1	0	-	1	-	-	-	-	-	1	-	-
	Cellular infiltration, inflammatory cell, renal pelvic mucosa	1	0	-	1	-	-	-	-	-	0	-	-
	Mineralization, renal pelvic mucosa	0	1	-	1	-	-	-	-	-	0	-	-
	Mineralization, cortex	0	0	-	0	-	-	-	-	-	1	-	-
	Uterine horn: Cellular infiltration, inflammatory cell	0	1	-	1	-	-	-	-	-	0	-	-
	Vagina: Atresia	0	2	-	2	-	-	-	-	-	0	-	-
	Pituitary gland: Cyst, pars intermedia	1	0	-	1	-	-	-	-	-	0	-	-
	Tubular hyperplasia, pars intermedia	1	0	-	1	-	-	-	-	-	1	-	-
	Mammary gland: Adenocarcinoma	1	0	-	1	-	-	-	-	-	1	-	-
F1	Number of animals examined	21	2	1	24 ^e	0	2	0	2	0	4	0	4
Findings in organs/tissues examined^{a,b,c}													
	Liver: Microgranuloma	11	1	0	12	-	1	-	1	-	3	-	3
	Kidney: Regeneration, tubular epithelium	1	0	-	1	-	0	-	0	-	1	-	1
	Dilatation, tubule	0	0	-	0	-	0	-	0	-	0	-	1
	Cellular infiltration, inflammatory cell, renal pelvic mucosa	1	0	-	1	-	0	-	0	-	0	-	0
	Cellular infiltration, inflammatory cell, cortex	0	0	-	0	-	0	-	0	-	0	-	1
	Mineralization, cortico-medullary junction	0	0	-	0	-	0	-	0	-	0	-	1
	Cast, hyaline	0	0	-	0	-	0	-	0	-	0	-	1
	Cyst	0	0	-	0	-	0	-	0	-	0	-	1
	Dilatation, renal pelvis	0	0	-	0	-	0	-	0	-	1	-	1
	Thymus: Atrophy, cortex	0	0	0	0	-	0	-	0	-	0	-	1
	Pituitary gland: Cyst, pars intermedia	1	0	0	1	-	0	-	0	-	1	-	0
	Tubular hyperplasia, pars intermedia	1	0	0	1	-	0	-	0	-	0	-	0
Findings in organs/tissues in which abnormal findings were grossly observed^{a,d}													
	Cerebrum: Dilatation, ventricle	-	-	-	-	-	-	-	-	-	-	1(1)	-
												-	1(1)

Fate: A, animals that had weanlings; B, animals that were not pregnant, did not produce viable pups or weanlings, or showed abnormal estrous cyclicity; C, animals that died during the study; T, total (A+B+C).

a: Values represent the number of animals that showed abnormal findings.

b: Organs/tissues examined were the liver, kidneys, thymus, thyroids, spleen, adrenals, bone marrow (femur), mesenteric lymph node, Peyer's patch, ovaries, uterus, vagina, mammary gland and pituitary gland.

c: Statistical analyses were made between the control and 4500 ppm groups based on the total number of animals examined.

d: Values in parentheses are the number of animals examined.

e: The number of animals examined on the kidney was 23 because both sides of the kidneys in one dead animal showed autolysis.

-: Not applicable.