

マダイ(Red sea-bream)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	18	100.0	18	100.0
0.01 ≤ C ≤ 0.1	0			
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

マハゼ (Spiny goby)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 ≤ C ≤ 0.1	0			
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ミナミマグロ (Southern bluefin tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.01	0.06	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	5	100.0	5	100.0
$0.1 < C \leq 0.2$	0			
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

メカジキ(Swordfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	0.02	0.04	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	5	100.0	5	100.0
$0.1 < C \leq 0.2$	0			
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

メバチ (Bigeye tuna)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	C<0.01	0.02	0.02

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	20.0	1	20.0
0.01 ≤ C ≤ 0.1	4	80.0	5	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

メバル(Brown rockfish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 ≤ C ≤ 0.1	0			
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ヤツメウナギ (Japanese lamprey)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	0.03	0.14	0.10

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	1	33.3	1	33.3
$0.1 < C \leq 0.2$	2	66.7	3	100.0
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

ヨシキリザメ (Great blue shark)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
5	C<0.01	0.05	0.03

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	20.0	1	20.0
0.01 ≤ C ≤ 0.1	4	80.0	5	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ワカサギ(Smelt)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
21	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	21	100.0	21	100.0
0.01 ≤ C ≤ 0.1	0			
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

2. 甲殻類

ガザミ(Common shoe-crab)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	C<0.01	0.29	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	8	26.7	8	26.7
0.01 ≤ C ≤ 0.1	15	50.0	23	76.7
0.1 < C ≤ 0.2	5	16.7	28	93.3
0.2 < C ≤ 0.3	2	6.7	30	100.0
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

クルマエビ (Kuruma-prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
35	C<0.01	0.41	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	17	48.6	17	48.6
0.01 ≤ C ≤ 0.1	15	42.9	32	91.4
0.1 < C ≤ 0.2	1	2.9	33	94.3
0.2 < C ≤ 0.3	1	2.9	34	97.1
0.3 < C ≤ 0.4	0	0.0	34	97.1
0.4 < C ≤ 0.5	1	2.9	35	100.0
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ケガニ (Korean crab (Muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	0.02	0.17	0.08

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	24	80.0	24	80.0
0.1 < C ≤ 0.2	6	20.0	30	100.0
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

スジエビ (Freshwater prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
18	0.02	0.11	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	17	94.4	17	94.4
$0.1 < C \leq 0.2$	1	5.6	18	100.0
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

テナガエビ (Scanpi)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	C<0.01	C<0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	3	100.0	3	100.0
0.01 ≤ C ≤ 0.1	0			
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ベニズワイガニ (Red snow crab)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	0.04	0.48	0.16

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	11	36.7	11	36.7
0.1 < C ≤ 0.2	9	30.0	20	66.7
0.2 < C ≤ 0.3	8	26.7	28	93.3
0.3 < C ≤ 0.4	1	3.3	29	96.7
0.4 < C ≤ 0.5	1	3.3	30	100.0
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ホッコクアカエビ (Deepwater prawn)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
45	0.02	0.57	0.11

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	33	73.3	33	73.3
$0.1 < C \leq 0.2$	5	11.1	38	84.4
$0.2 < C \leq 0.3$	2	4.4	40	88.9
$0.3 < C \leq 0.4$	4	8.9	44	97.8
$0.4 < C \leq 0.5$	0	0.0	44	97.8
$0.5 < C \leq 0.6$	1	2.2	45	100.0
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

3. 軟体動物

アカガイ (Chest-shell)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	0.60	0.68	0.64

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	0	0.0	0	0.0
0.1 < C ≤ 0.2	0	0.0	0	0.0
0.2 < C ≤ 0.3	0	0.0	0	0.0
0.3 < C ≤ 0.4	0	0.0	0	0.0
0.4 < C ≤ 0.5	0	0.0	0	0.0
0.5 < C ≤ 0.6	1	33.3	1	33.3
0.6 < C ≤ 0.7	2	66.7	3	100.0
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

アサリ (Japanese short-neck)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
51	0.02	0.17	0.06

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	49	96.1	49	96.1
$0.1 < C \leq 0.2$	2	3.9	51	100.0
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

アワビ (Abalone(muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	0.02	0.07	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	15	100.0	15	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

イイダコ(Ocellated octopus)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	0.04	0.05	0.04

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	3	100.0	3	100.0
$0.1 < C \leq 0.2$	0			
$0.2 < C \leq 0.3$	0			
$0.3 < C \leq 0.4$	0			
$0.4 < C \leq 0.5$	0			
$0.5 < C \leq 0.6$	0			
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

コウイカ (Cuttlefish)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	1	6.7	1	6.7
0.01 ≤ C ≤ 0.1	14	93.3	15	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

サザエ (Horned turban (Muscle))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	0.10	0.05

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	2	13.3	2	13.3
0.01 ≤ C ≤ 0.1	13	86.7	15	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

シジミ (Japanese corbicula)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
64	0.03	0.77	0.37

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	7	10.9	7	10.9
0.1 < C ≤ 0.2	11	17.2	18	28.1
0.2 < C ≤ 0.3	7	10.9	25	39.1
0.3 < C ≤ 0.4	21	32.8	46	71.9
0.4 < C ≤ 0.5	0	0.0	46	71.9
0.5 < C ≤ 0.6	0	0.0	46	71.9
0.6 < C ≤ 0.7	9	14.1	55	85.9
0.7 < C ≤ 0.8	9	14.1	64	100.0
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

スルメイカ (Pacific flying squid)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
56	0.03	1.30	0.29

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	18	32.1	18	32.1
0.1 < C ≤ 0.2	11	19.6	29	51.8
0.2 < C ≤ 0.3	9	16.1	38	67.9
0.3 < C ≤ 0.4	6	10.7	44	78.6
0.4 < C ≤ 0.5	4	7.1	48	85.7
0.5 < C ≤ 0.6	1	1.8	49	87.5
0.6 < C ≤ 0.7	1	1.8	50	89.3
0.7 < C ≤ 0.8	1	1.8	51	91.1
0.8 < C ≤ 0.9	0	0.0	51	91.1
0.9 < C ≤ 1.0	2	3.6	53	94.6
1.0 < C ≤ 1.1	2	3.6	55	98.2
1.1 < C ≤ 1.2	0	0.0	55	98.2
1.2 < C ≤ 1.3	1	1.8	56	100.0
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ハマグリ (Japanese hard clam)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
48	0.02	0.14	0.07

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	40	83.3	40	83.3
0.1 < C ≤ 0.2	8	16.7	48	100.0
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ホタテ貝柱 (Common scallop (Muskelsehne))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
57	0.01	0.51	0.12

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	42	73.7	42	73.7
$0.1 < C \leq 0.2$	7	12.3	49	86.0
$0.2 < C \leq 0.3$	1	1.8	50	87.7
$0.3 < C \leq 0.4$	3	5.3	53	93.0
$0.4 < C \leq 0.5$	2	3.5	55	96.5
$0.5 < C \leq 0.6$	2	3.5	57	100.0
$0.6 < C \leq 0.7$	0			
$0.7 < C \leq 0.8$	0			
$0.8 < C \leq 0.9$	0			
$0.9 < C \leq 1.0$	0			
$1.0 < C \leq 1.1$	0			
$1.1 < C \leq 1.2$	0			
$1.2 < C \leq 1.3$	0			
$1.3 < C \leq 1.4$	0			
$1.4 < C \leq 1.5$	0			
$1.5 < C \leq 1.6$	0			
$1.6 < C \leq 1.7$	0			
$1.7 < C \leq 1.8$	0			
$1.8 < C \leq 1.9$	0			
$1.9 < C \leq 2.0$	0			
$2.0 < C \leq 2.1$	0			
$2.1 < C \leq 2.2$	0			
$2.2 < C \leq 2.3$	0			
$2.3 < C \leq 2.4$	0			
$2.4 < C \leq 2.5$	0			
$2.5 < C \leq 2.6$	0			
$2.6 < C \leq 2.7$	0			
$2.7 < C \leq 2.8$	0			
$2.8 < C \leq 2.9$	0			
$2.9 < C \leq 3.0$	0			
$3.0 < C$	0			

マガキ (Giant pacific oyster)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
45	0.10	0.68	0.30

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	1	2.2	1	2.2
0.1 < C ≤ 0.2	16	35.6	17	37.8
0.2 < C ≤ 0.3	8	17.8	25	55.6
0.3 < C ≤ 0.4	7	15.6	32	71.1
0.4 < C ≤ 0.5	8	17.8	40	88.9
0.5 < C ≤ 0.6	2	4.4	42	93.3
0.6 < C ≤ 0.7	3	6.7	45	100.0
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

マダコ(Octopus)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
24	C<0.01	0.07	0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	14	58.3	14	58.3
0.01 ≤ C ≤ 0.1	10	41.7	24	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

4. 棘皮動物
ウニ (Sea urchin)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
45	0.02	0.34	0.17

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	8	17.8	8	17.8
0.1 < C ≤ 0.2	24	53.3	32	71.1
0.2 < C ≤ 0.3	10	22.2	42	93.3
0.3 < C ≤ 0.4	3	6.7	45	100.0
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

ナマコ (Sea cucumber)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	C<0.01	0.01	C<0.01

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	14	93.3	14	93.3
0.01 ≤ C ≤ 0.1	1	6.7	15	100.0
0.1 < C ≤ 0.2	0			
0.2 < C ≤ 0.3	0			
0.3 < C ≤ 0.4	0			
0.4 < C ≤ 0.5	0			
0.5 < C ≤ 0.6	0			
0.6 < C ≤ 0.7	0			
0.7 < C ≤ 0.8	0			
0.8 < C ≤ 0.9	0			
0.9 < C ≤ 1.0	0			
1.0 < C ≤ 1.1	0			
1.1 < C ≤ 1.2	0			
1.2 < C ≤ 1.3	0			
1.3 < C ≤ 1.4	0			
1.4 < C ≤ 1.5	0			
1.5 < C ≤ 1.6	0			
1.6 < C ≤ 1.7	0			
1.7 < C ≤ 1.8	0			
1.8 < C ≤ 1.9	0			
1.9 < C ≤ 2.0	0			
2.0 < C ≤ 2.1	0			
2.1 < C ≤ 2.2	0			
2.2 < C ≤ 2.3	0			
2.3 < C ≤ 2.4	0			
2.4 < C ≤ 2.5	0			
2.5 < C ≤ 2.6	0			
2.6 < C ≤ 2.7	0			
2.7 < C ≤ 2.8	0			
2.8 < C ≤ 2.9	0			
2.9 < C ≤ 3.0	0			
3.0 < C	0			

5. 魚介類内臓

アカニシ(内臓含む)(Thomas,s rapa whelk)

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
3	1.50	3.80	2.53

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.1	0	0.0	0	0.0
0.1 < C ≤ 0.2	0	0.0	0	0.0
0.2 < C ≤ 0.3	0	0.0	0	0.0
0.3 < C ≤ 0.4	0	0.0	0	0.0
0.4 < C ≤ 0.5	0	0.0	0	0.0
0.5 < C ≤ 0.6	0	0.0	0	0.0
0.6 < C ≤ 0.7	0	0.0	0	0.0
0.7 < C ≤ 0.8	0	0.0	0	0.0
0.8 < C ≤ 0.9	0	0.0	0	0.0
0.9 < C ≤ 1.0	0	0.0	0	0.0
1.0 < C ≤ 1.1	0	0.0	0	0.0
1.1 < C ≤ 1.2	0	0.0	0	0.0
1.2 < C ≤ 1.3	0	0.0	0	0.0
1.3 < C ≤ 1.4	1	33.3	1	33.3
1.4 < C ≤ 1.5	0	0.0	1	33.3
1.5 < C ≤ 1.6	0	0.0	1	33.3
1.6 < C ≤ 1.7	0	0.0	1	33.3
1.7 < C ≤ 1.8	0	0.0	1	33.3
1.8 < C ≤ 1.9	0	0.0	1	33.3
1.9 < C ≤ 2.0	0	0.0	1	33.3
2.0 < C ≤ 2.1	0	0.0	1	33.3
2.1 < C ≤ 2.2	0	0.0	1	33.3
2.2 < C ≤ 2.3	1	33.3	2	66.7
2.3 < C ≤ 2.4	0	0.0	2	66.7
2.4 < C ≤ 2.5	0	0.0	2	66.7
2.5 < C ≤ 2.6	0	0.0	2	66.7
2.6 < C ≤ 2.7	0	0.0	2	66.7
2.7 < C ≤ 2.8	0	0.0	2	66.7
2.8 < C ≤ 2.9	0	0.0	2	66.7
2.9 < C ≤ 3.0	0	0.0	2	66.7
3.0 < C	1	33.3	3	100.0

アワビ内臓 (Abalone (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	2.20	5.60	3.11

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.5$	0	0.0	0	0.0
$0.5 < C \leq 1.0$	0	0.0	0	0.0
$1.0 < C \leq 1.5$	0	0.0	0	0.0
$1.5 < C \leq 2.0$	0	0.0	0	0.0
$2.0 < C \leq 2.5$	1	6.7	1	6.7
$2.5 < C \leq 3.0$	8	53.3	9	60.0
$3.0 < C \leq 3.5$	4	26.7	13	86.7
$3.5 < C \leq 4.0$	1	6.7	14	93.3
$4.0 < C \leq 4.5$	0	0.0	14	93.3
$4.5 < C \leq 5.0$	0	0.0	14	93.3
$5.0 < C \leq 5.5$	0	0.0	14	93.3
$5.5 < C \leq 6.0$	1	6.7	15	100.0
$6.0 < C \leq 6.5$	0			
$6.5 < C \leq 7.0$	0			
$7.0 < C \leq 7.5$	0			
$7.5 < C \leq 8.0$	0			
$8.0 < C \leq 8.5$	0			
$8.5 < C \leq 9.0$	0			
$9.0 < C \leq 9.5$	0			
$9.5 < C \leq 10$	0			
$10 < C$	0			

ガザミみそ(Common shoe-crab (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
30	0.09	1.90	0.69

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
$C < 0.01$	0	0.0	0	0.0
$0.01 \leq C \leq 0.1$	2	6.7	2	6.7
$0.1 < C \leq 0.2$	6	20.0	8	26.7
$0.2 < C \leq 0.3$	4	13.3	12	40.0
$0.3 < C \leq 0.4$	2	6.7	14	46.7
$0.4 < C \leq 0.5$	0	0.0	14	46.7
$0.5 < C \leq 0.6$	0	0.0	14	46.7
$0.6 < C \leq 0.7$	0	0.0	14	46.7
$0.7 < C \leq 0.8$	3	10.0	17	56.7
$0.8 < C \leq 0.9$	2	6.7	19	63.3
$0.9 < C \leq 1.0$	5	16.7	24	80.0
$1.0 < C \leq 1.1$	0	0.0	24	80.0
$1.1 < C \leq 1.2$	0	0.0	24	80.0
$1.2 < C \leq 1.3$	3	10.0	27	90.0
$1.3 < C \leq 1.4$	0	0.0	27	90.0
$1.4 < C \leq 1.5$	1	3.3	28	93.3
$1.5 < C \leq 1.6$	1	3.3	29	96.7
$1.6 < C \leq 1.7$	0	0.0	29	96.7
$1.7 < C \leq 1.8$	0	0.0	29	96.7
$1.8 < C \leq 1.9$	1	3.3	30	100.0
$1.9 < C \leq 2.0$	0			
$2.0 < C$	0			

ケガニみそ (Korean crab (internal organ))

全試料数	最小値 (mg/kg)	最大値 (mg/kg)	平均値 (mg/kg)
15	0.79	3.50	1.99

カドミウム濃度 (mg/kg)	試料数	割合 (%)	累積試料数	割合 (%)
C < 0.01	0	0.0	0	0.0
0.01 ≤ C ≤ 0.5	0	0.0	0	0.0
0.5 < C ≤ 1.0	1	6.7	1	6.7
1.0 < C ≤ 1.5	4	26.7	5	33.3
1.5 < C ≤ 2.0	2	13.3	7	46.7
2.0 < C ≤ 2.5	5	33.3	12	80.0
2.5 < C ≤ 3.0	2	13.3	14	93.3
3.0 < C ≤ 3.5	1	6.7	15	100.0
3.5 < C ≤ 4.0	0			
4.0 < C ≤ 4.5	0			
4.5 < C ≤ 5.0	0			
5.0 < C ≤ 5.5	0			
5.5 < C ≤ 6.0	0			
6.0 < C ≤ 6.5	0			
6.5 < C ≤ 7.0	0			
7.0 < C ≤ 7.5	0			
7.5 < C ≤ 8.0	0			
8.0 < C ≤ 8.5	0			
8.5 < C ≤ 9.0	0			
9.0 < C ≤ 9.5	0			
9.5 < C ≤ 10	0			
10 < C	0			